

Lupaneta Pack™

leuprolide acetate for depot suspension
and norethindrone acetate tablets

Treat your endometriosis. Take care of you.



Use

What is LUPANETA PACK™ (leuprolide acetate for depot suspension and norethindrone acetate tablets)?

LUPANETA PACK is used for initial management of the painful symptoms of endometriosis and for management of recurrence of symptoms. It contains 2 different prescription medicines:

- Leuprolide acetate for depot suspension, a medicine injected into your muscle and used to treat pain due to endometriosis
- Norethindrone acetate tablets, a medicine taken by mouth and used to help lower the side effect of bone thinning that is caused by leuprolide acetate for depot suspension

LUPANETA PACK should not be used longer than 6 months at a time after you first start treatment for your endometriosis symptoms. LUPANETA PACK should not be used for more than a total of 12 months during your treatment.

It is not known if LUPANETA PACK is safe and effective in children under 18 years of age.

Safety Considerations

- Do not take LUPANETA PACK™ if you are or may become pregnant; are breast-feeding; have undiagnosed uterine bleeding; a history of breast or other cancers; problems with blood clots; or stroke, heart attack, or liver problems
- Serious side effects may include thinning of bones which may not be completely reversible, vision problems (including sudden loss of vision, double vision, bulging eyes, or migraine headaches), depression or worsening of depression, and swelling (fluid retention)
- A condom, a diaphragm with contraceptive jelly, or a copper IUD is required to prevent pregnancy
- Increased endometriosis symptoms (such as pelvic pain or pressure and/or pain during intercourse) may occur for 1 or 2 weeks after starting therapy
- Reports of convulsions have occurred in patients taking leuprolide acetate

Please see Use and additional Important Safety Information on back cover.

Please see accompanying full Prescribing Information, here:

[LUPANETA PACK Prescribing Information.](#)

Treat your endometriosis.


Take care of you.

LUPANETA PACK™

Relieves Endometriosis Pain

Leuprolide acetate for depot suspension helps manage endometriosis pain. It comes as an injection that your healthcare provider administers.¹

In a Clinical Study,
Women Experienced a
Significant Reduction in Painful Symptoms¹



Before treatment with leuprolide acetate and norethindrone acetate tablets

After treatment with leuprolide acetate and norethindrone acetate tablets

The charts above show the percentages of women in a clinical study who had certain types of endometriosis pain before treatment and after about a year of LUPANETA PACK therapy. Women's symptoms of painful periods and pelvic pain were evaluated in 55 women; symptoms of pain during intercourse were evaluated in 43 women.¹

Lupaneta Pack™

leuprolide acetate for depot suspension
and norethindrone acetate tablets

Learn more at LUPANETAPACK.COM

Please see Use and Important Safety Information on back cover.
Please see accompanying full Prescribing Information, here:
[LUPANETA PACK Prescribing Information.](#)

LUPANETA PACK™

Reduces Hot Flashes & Bone Loss

Norethindrone acetate tablets (add-back) add a small amount of progestin to help with hot flashes and bone loss associated with leuprolide acetate. In one study, after 6 months of therapy, women taking LUPANETA PACK reported¹:

63%

fewer average
number of days
with hot flashes


than women taking
leuprolide acetate alone

67%

fewer average
maximum number of
hot flashes per day

than women taking
leuprolide acetate alone

Women taking LUPANETA PACK experienced significantly less bone density loss than women taking leuprolide acetate for depot suspension alone¹:



Safety Considerations

The most common side effects of LUPANETA PACK include hot flashes and sweats, headaches or migraine headaches, depression and mood swings, nausea and vomiting, problems sleeping, and nervousness or feeling anxious. Tell your doctor if you have any side effect that bothers you or that does not go away.¹

LUPANETA PACK™ can help.

LUPANETA PACK relieves endometriosis pain and reduces hot flashes and bone loss.

Leuprolide acetate for depot suspension injection suppresses estrogen production, creating a low-estrogen state that can reduce endometriosis pain.¹


Norethindrone acetate 5 mg tablets (also called “add-back”) add a small amount of progestin to help reduce hot flashes and bone loss associated with leuprolide acetate—without reducing how well your treatment works.¹

What to expect during a full 6-month course of LUPANETA PACK therapy¹

Estrogen levels temporarily increase after treatment

Increased estrogen levels may temporarily worsen your symptoms

Estrogen levels decrease after 1 to 2 weeks, and you start to feel relief from endometriosis pain



For illustrative purposes only. Responses may vary by individual patients.

If your endometriosis symptoms recur after your initial course of therapy, your doctor can prescribe an additional 6 months of treatment. Your doctor should assess your bone density before retreatment. Retreatment should be limited to 6 months.¹

LUPANETA PACK **does NOT** put you into menopause, although you may experience similar effects such as hot flashes and a break from your normal menstrual cycle. These effects are **temporary** and **reversible** after therapy is stopped.¹

In a clinical study, **95% of women resumed a normal cycle within 3 months after the end of LUPANETA PACK therapy with leuprolide acetate.**⁴

Safety Considerations

Do not take LUPANETA PACK if you:

- Have had an allergic reaction to medicines like leuprolide acetate or norethindrone acetate tablets
- Have uterine bleeding for which a cause has not been found
- Are pregnant or may be pregnant
- Are breast-feeding or plan to breast-feed
- Had or have breast cancer or other cancers that are sensitive to hormones
- Have problems with blood clots, a stroke, or a heart attack
- Have liver problems

Lupaneta Pack™

leuprolide acetate for depot suspension
and norethindrone acetate tablets


Please see Use and Important Safety Information on back cover. Please see accompanying full Prescribing Information, here: [LUPANETA PACK Prescribing Information](#).

You've probably been dealing with endometriosis symptoms for years.

Taking control can start here.

Endometriosis is a common health problem in women. It occurs when endometrium—tissue that normally lines the uterus—grows outside the uterus. These abnormal growths are called lesions.²

Estrogen fuels the growth of lesions, which then bleed and break down, causing your painful symptoms.²



To download your LUPANETA PACK Savings Card* now visit:

LupanetaPack.com

If you have additional questions regarding savings and support, call: 1-855-587-7663

***Eligibility:** Available to patients in the United States with commercial prescription insurance coverage for LUPANETA PACK™ (leuprolide acetate for depot suspension and norethindrone acetate tablets) who meet eligibility criteria. Co-pay assistance program is not available to patients receiving prescription reimbursement under any federal, state or government-funded insurance programs (for example, Medicare (including Part D), Medicare Advantage, Medigap, Medicaid, TRICARE, Department of Defense or Veteran's Affairs programs) or where prohibited by law or by the patient's health insurance provider. If at any time a patient begins receiving prescription drug coverage under any such federal, state or government-funded healthcare program, patient will no longer be able to use the LUPRON DEPOT® card and patient must call 1-855-587-7663 to stop participation. Patients residing in or receiving treatment in certain states may not be eligible. Patients may not seek reimbursement for value received from this card from any third-party payers. Offer subject to change or discontinuance without notice. Restrictions, including monthly maximums, may apply. **This is not health insurance.**

Good for out-of-pocket expenses up to \$125 on your 3.75 mg LUPANETA PACK prescription after the initial \$10 co-pay or up to \$250 on your – 3 Month 11.25 mg LUPANETA PACK prescription after the initial \$10 co-pay and up to \$25 for generic norethindrone acetate after the initial \$5 co-pay on six occasions when accompanied with a prescription for LUPANETA PACK. AbbVie reserves the right to rescind, revoke, or amend this offer without notice. LUPANETA PACK is a registered trademark of AbbVie Inc. You may not combine this offer with any other rebate, coupon, free trial, or similar offer.

References

1. LUPANETA PACK [package insert]. 2. The American College of Obstetricians and Gynecologists (ACOG). FAQ Endometriosis. ACOG website. <http://www.acog.org/Patients/FAQs/Endometriosis>. Published October 2012. Accessed April 30, 2015. 3. Office on Women's Health. Endometriosis. Womenshealth.gov website. <http://womenshealth.gov/publications/our-publications/fact-sheet/endometriosis.html>. Published August 18, 2014. Accessed April 30, 2015. 4. LUPRON DEPOT [package insert].

You are not alone.

About 5 million women
in the United States suffer with endometriosis.³

Treat your endometriosis. Take care of you.

Learn more at LUPANETAPACK.COM

Lupaneta Pack™
leuprolide acetate for depot suspension
and norethindrone acetate tablets

Use

What is LUPANETA PACK™ (leuprolide acetate for depot suspension and norethindrone acetate tablets)?

LUPANETA PACK is used for initial management of the painful symptoms of endometriosis and for management of recurrence of symptoms. It contains 2 different prescription medicines:

- Leuprolide acetate for depot suspension, a medicine injected into your muscle and used to treat pain due to endometriosis
- Norethindrone acetate tablets, a medicine taken by mouth and used to help lower the side effect of bone thinning that is caused by leuprolide acetate for depot suspension

LUPANETA PACK should not be used longer than 6 months at a time after you first start treatment for your endometriosis symptoms. LUPANETA PACK should not be used for more than a total of 12 months during your treatment.

It is not known if LUPANETA PACK is safe and effective in children under 18 years of age.

Important Safety Information

Who should not take LUPANETA PACK?

Do not take LUPANETA PACK if you:

- Have had an allergic reaction to medicines like leuprolide acetate or norethindrone acetate tablets
- Have uterine bleeding for which a cause has not been found
- Are pregnant or may be pregnant
- Are breast-feeding or plan to breast-feed
- Had or have breast cancer or other cancers that are sensitive to hormones
- Have problems with blood clots, a stroke, or a heart attack
- Have liver problems

What should I tell my doctor before taking LUPANETA PACK?

Before you take LUPANETA PACK, tell your doctor if you:

- Drink alcohol
- Smoke
- Have a family history of bone loss (osteoporosis)
- Have or have had depression
- Have high cholesterol
- Have had blood clots, a stroke, or a heart attack
- Have migraine headaches
- Have diabetes
- Have epilepsy
- Have kidney problems

Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements.

Especially tell your doctor if you take anticonvulsant (seizure) or corticosteroid medicines.

Before you take LUPANETA PACK, talk to your doctor about a birth control method that is right for you. You will need to use a form of birth control that does not contain hormones, such as:

- A diaphragm with spermicide
- Condoms with spermicide
- A copper IUD

If you become pregnant while taking LUPANETA PACK, stop taking the norethindrone acetate tablets and call your doctor right away.

What are the possible side effects of LUPANETA PACK?

LUPANETA PACK may cause serious side effects, including:

- Bone thinning (decreased bone mineral density)
 - Harm to your unborn baby
 - Vision problems. Call your doctor right away if you have sudden loss of vision, double vision, bulging eyes, or migraine headaches
 - Depression or worsening depression
 - Worsening endometriosis symptoms when you start taking LUPANETA PACK
 - Swelling (fluid retention)
 - Convulsions
 - Allergic reactions.
- Get medical help right away if you have any of these symptoms: – trouble breathing – swelling of your face, lips, mouth, or tongue – wheezing – severe itching – skin rash, redness, or swelling – dizziness or fainting – fast heartbeat or pounding in your chest (tachycardia) – sweating

The most common side effects of LUPANETA PACK include:

- Hot flashes and sweats
- Headaches or migraine headaches
- Depression and mood swings
- Nausea and vomiting
- Problems sleeping
- Nervousness or feeling anxious
- Pain
- Acne
- Weakness
- Vaginal infection or inflammation
- Weight gain
- Constipation or diarrhea

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all of the possible side effects of LUPANETA PACK. For more information, ask your doctor or pharmacist.

Please see accompanying full Prescribing Information, here:

[LUPANETA PACK Prescribing Information.](#)